arXiv:0802.3745v2 [math.DS] 26 Sep 2009

HETERODIMENSIONAL TANGENCIES ON CYCLES LEADING TO STRANGE ATTRACTORS

SHIN KIRIKI, YUSUKE NISHIZAWA, AND TERUHIKO SOMA

Abstract. In this paper, we study a two-parameter family $\{\varphi_{\mu,\nu}\}$ of 3-dimensional diffeomorphisms which have a bifurcation induced by simultaneous generation of a heterodimensional cycle and a heterodimensional tangency associated to two saddle points. We show that such a codimension-2 bifurcation generates a quadratic homoclinic tangency associated to one of the saddle continuations which unfolds generically with respect to some one-parameter subfamily of $\{\varphi_{\mu,\nu}\}$. Moreover, from this result together with some well-known facts, we detect some nonhyperbolic phenomena (i.e., the existence of nonhyperbolic strange attractors and the C^2 robust tangencies) arbitrarily close to the codimension-2 bifurcation.

1. Introduction

When diffeomorphisms act on manifolds of dimension greater than or equal to three, it is well known that nonhyperbolic phenomena are caused by the existence of heteroclinic cycles containing two saddle points with different indexes, called *heterodimensional cycles*, as well as that of homoclinic tangencies [13, 16, 10, 15, 17, 11]. Heterodimensional cycles presented a new mechanism in dynamics, which has been studied Díaz et al. [7, 5, 8, 2, 9, 3, 4]. As is suggested in [14], these two classes of nonhyperbolic diffeomorphisms seem to occupy a large part in the complement of the hyperbolic diffeomorphisms. In this paper, we study 3-dimensional diffeomorphisms which have heterodimensional cycles and tangencies of certain type simultaneously.

Let φ be a diffeomorphism on a 3-dimensional smooth manifold M which has two saddle fixed points p and q satisfying $\operatorname{index}(q) = \operatorname{index}(p) + 1$, where $\operatorname{index}(\cdot)$ denotes the dimension of the unstable manifold of the saddle point. A heteroclinic point r of the stable manifold $W^s(p)$ and the unstable manifold $W^u(q)$ is called a *heterodimensional tangency* of $W^s(p)$ and $W^u(q)$ if r satisfies

- $T_rW^s(p) + T_rW^u(q) \neq T_rM$;
- $\dim(T_rW^s(p)) + \dim(T_rW^u(q)) > \dim(M)$,

where $\dim(\cdot)$ denotes the dimension of the space.

Our main theorem in this paper is as follows. We will present some definitions and generic conditions used in the statement of Theorem A in Section 2.

Theorem A. *Let M be a 3-dimensional C^2 manifold, and let $\{\varphi_{\mu,\nu}\}$ be a two-parameter family of C^2 diffeomorphisms $\varphi_{\mu,\nu} : M \to M$ which C^2 depends on (μ, ν)*

Date: September 26, 2009, to appear in *Discrete Conti. Dynam. Sys.*

2000 *Mathematics Subject Classification.* Primary:37C29, 37D45, 37G30.

Key words and phrases. heterodimensional cycle, homoclinic tangency, strange attractors, infinitely many sinks.

and has continuations of saddle fixed points $p_{\mu,\nu}$ *and* $q_{\mu,\nu}$ *with* $\mathrm{index}(p_{\mu,\nu}) = 1$ *and* $\mathrm{index}(q_{\mu,\nu}) = 2$*. Suppose that the following conditions hold.*

- *Each* $\varphi_{\mu,\nu}$ *is locally* C^2 *linearizable in a small neighborhood* $N(q_{\mu,\nu})$ *of* $q_{\mu,\nu}$*.*
- $\varphi = \varphi_{0,0}$ *has a heterodimensional cycle containing the fixed points* $p = p_{0,0}$*,* $q = q_{0,0}$*, a nondegenerate heterodimensional tangency* r*, a quasi-transverse intersection* $s \in W^s(q) \cap W^u(p)$*.*
- $\{\varphi_{\mu,\nu}\}$ *satisfies the generic conditions (C1)-(C4) given in Section 2.*

Then, for a sufficiently small $\varepsilon > 0$ *and any* μ *in either* $(0, \varepsilon)$ *or* $(-\varepsilon, 0)$*, there exist infinitely many* ν *such that* $\varphi_{\mu,\nu}$ *has a quadratic homoclinic tangency associated to* $p_{\mu,\nu}$ *which unfolds generically with respect to the* ν*-parameter family* $\{\varphi_{\mu(\mathrm{fixed}),\nu}\}$*.*

Remark 1.1. (1) The generic conditions (C2) and (C3) imply the setting such that a heterodimensional tangency and a quasi-transverse intersection unfold generically with respect to the parameters μ and ν, respectively. The point to notice is that the newly detected homoclinic tangency in Theorem A can be also controlled by these parameters. It is not hard to get a one parameter family in the infinite dimensional space $\mathrm{Diff}^2(M)$ with respect to which the homoclinic tangency unfolds generically. However, in our proof, we need to show that the tangency given in Subsection 4.1 unfolds generically with respect to the ν-parameter family $\{\varphi_{\mu_0,\nu}\}$ in the two-dimensional subspace $\{\varphi_{\mu,\nu}\}$ of $\mathrm{Diff}^2(M)$.

(2) Theorem A holds for a homoclinic tangency associated to $q_{\mu,\nu}$ instead of $p_{\mu,\nu}$ if we replace the conditions in (C1) and (C4) by appropriate ones.

(3) We need the C^2 smoothness in the local linearization around $q_{\mu,\nu}$ to estimate the curvatures of $W^s(p_{\mu,\nu})$ and $W^u(p_{\mu,\nu})$ at the tangency in Section 3.

Figure 1.1 illustrates an example of heterodimensional cycles containing a nondegenerate heterodimensional tangency of hyperbolic type, see Definition 2.1-(3).



Figure 1.1. A heterodimensional cycle associated with the pair of saddles p and q, which contains the heterodimensional tangency r and the quasi-transverse intersection s.

The conclusion obtained from Theorem A reminds us of prior works associated with homoclinic tangencies. The one is related to strange attractors and the other C^2 robust tangencies.

First, let us discuss the former one. Viana showed the following theorem.

Theorem 1.2 (Viana [18]). *For a generic subset of one-parameter families* $\{\varphi_\mu\}$ *of* C^∞ *diffeomorphisms on any manifolds of the dimension greater than or equal to two that unfolds a homoclinic tangency at parameter value* $\mu = 0$ *associated to a sectionally dissipative saddle periodic point, there is a subset* S *of* $\mathbb{R}$ *such that*

- $S \cap (-\epsilon, \epsilon)$ *has a positive Lebesgue measure for every* $\epsilon > 0$,
- *for all* $\mu \in S$, φ_μ *exhibits nonhyperbolic strange attractors in a* μ*-dependent neighborhood of the orbit of tangency.*

Leal [12] extended this result and showed the existence of infinitely many strange attractors. A saddle periodic point is said to be *sectionally dissipative* if the product of any two eigenvalues of the derivative at the point has norm less than one. Also, Λ is a *strange attractor* of φ if Λ is a compact, φ-invariant, transitive set and the basin $W^s(\Lambda)$ has a nonempty interior and there exists $z_1 \in \Lambda$ such that $\{\varphi^n(z_1) : n \geq 0\}$ is dense in Λ and $||d\varphi^n(z_1)|| \geq e^{cn}$ for all $n \geq 0$ and some $c > 0$. Note, Viana [18] assumed that for simplicity φ_μ is C^3 linearizable in a neighborhood of the saddle point for μ is sufficiently close to 0. Combining these extra conditions and our result for the cycle containing the heterodimensional tangency imply the following corollary.

Corollary B. *Let* $\{\varphi_{\mu,\nu}\}$ *be the two-parameter family of* 3*-dimensional* C^∞ *diffeomorphisms having the heterodimensional cycle with the nondegenerate heterodimensional tangency for* $(\mu, \nu) = (0, 0)$ *given in Theorem A. Suppose that each* $\varphi_{\mu,\nu}$ *is locally* C^3 *linearizable in a small neighborhood of* $p_{\mu,\nu}$ *and* $\varphi_{0,0}$ *is sectionally dissipative at* $p_{0,0}$. *Then there exists a positive Lebesgue measure subset* $\mathcal{A}$ *of the* $\mu\nu$*-plane arbitrarily near* $(0, 0)$ *such that, for any* $(\mu, \nu) \in \mathcal{A}$, $\varphi_{\mu,\nu}$ *exhibits nonhyperbolic strange attractors.*

Next, we discuss C^2 robust homoclinic tangencies derived from the heterodimensional cycle having a nondegenerate tangency. A homoclinic tangency of a diffeomorphism φ associated with a hyperbolic set Γ is C^r *robust* if there is a C^r neighborhood $\mathcal{U}$ of φ such that every diffeomorphism $\psi \in \mathcal{U}$ has a homoclinic tangency associated with the continuations of Γ for ψ. Newhouse [13] showed, in the C^2 topology, a homoclinic tangency of surface diffeomorphisms generates C^2 robust homoclinic tangencies. This property yields the so-called C^2 Newhouse phenomenon: there is a non-empty open set of C^2 diffeomorphisms and its residual subset such that every diffeomorphism in the subset has infinitely many sinks. The result is extended by Palis-Viana [15] to the higher dimensional case. Palis and Viana proved the result under the sectional dissipativeness and linearizing conditions as in [18]. Moreover, Romero [17] proved the following theorem without these conditions.

Theorem 1.3 (Romero [17]). *Let* φ *be a* C^2 *diffeomorphism on a manifold* M, $\dim(M) \geq 3$, *having a homoclinic tangency associated with a saddle periodic point of* φ *whose index is greater or equal to* 2. *Then there are diffeomorphisms arbitrarily* C^2 *close to* φ *having robust homoclinic tangencies.*

Therefore, we have the following corollary.

Corollary C. *Let $\{\varphi_{\mu,\nu}\}$ be the two-parameter family of 3-dimensional C^2 diffeomorphisms having the heterodimensional cycle with the nondegenerate heterodimensional tangency for $(\mu,\nu)=(0,0)$ given in Theorem A. Then, for a sufficiently small $\varepsilon>0$ and any μ in either $(0,\varepsilon)$ or $(-\varepsilon,0)$, there are infinitely many ν such that every C^2 neighborhood of $\varphi_{\mu,\nu}$ contains a diffeomorphism having a C^2 robust homoclinic tangency.*

Remark 1.4. If we add a *weak* dissipative condition (see in [17, Theorem A (1.1)]) to the assumptions of Corollary C then, the C^2 Newhouse phenomenon is obtained from our settings.

Remark 1.5. In the C^1 case, Díaz et al. [6] have already proved that the unfolding of heterodimensional tangencies leads to non-dominated dynamics and therefore (by results of [2] and [3]) to the C^1 Newhouse phenomenon (see also [1] for a different approach to the phenomenon). On the other hand, the theory of strange attractors has not been so far developed in the C^l category ($l=1,2$).

Outline of the proof of Theorem A: We will finish Introduction by presenting a sketch of the proof of Theorem A.

In Section 2, we give definitions and the generic conditions (C1)-(C4) used in Theorem A. Especially, a nondegenerate heterodimensional tangency, which is one of main ingredients of this paper, is introduced explicitly there. Such tangencies are classified into the elliptic and hyperbolic types, see Definition 2.1-(3).



FIGURE 1.2. (1) The situation when $(\mu,\nu)=(0,0)$. (2) $s_{\mu,0}$ is a quasi-transverse intersection with respect to the new parameter. (3) l_m converges $W^{uu}_{\mathrm{loc}}(q_{\mu,0})$ in C^2 topology. (4) l_{m,ν_m} is a continuation of l_m for $\nu=\nu_m$.

In Section 3, we prove three lemmas. Lemma 3.1 shows the existence of the reparametrization of $\{\varphi_{\mu,\nu}\}$ such that, for any $(\mu,0)$ in the new parameter, $W^s(q_{\mu,0})$

and $W^u(p_{u,0})$ still have a quasi-transverse intersection $s_{\mu,0}$ which unfolds generically with respect to the ν-parameter but the tangency r is annihilated. See Fig.1.2 (1) and (2). Lemma 3.2 applies the C^2 inclination lemma to a shorter curve l_{m_0} in $W^u(p_{\mu,0})$ passing through $s_{\mu,0}$ so that some curve l_m in $\varphi^m_{\mu,0}(l_{m_0})$ containing $\varphi^m_{\mu,0}(s_{\mu,0})$ C^2 converges to $W^{uu}_{\mathrm{loc}}(q_{\mu,0})$ as $m \to \infty$. See Fig.1.2 (3). Lemma 3.3 explains a connection between quadratic tangencies and curvatures, which is used to show that the homoclinic tangencies obtained in Section 4 are quadratic.

Assertion 4.2 and Assertion 4.3 in Section 4 show that the generic unfolding of heterodimensional tangencies introduces the existence of a quadratic homoclinic tangency τ_m as illustrated in Fig. 1.2-(4). Finally, we prove in Proposition 4.4 that the tangency τ_m unfolds generically with respect to the ν-parameter. These results assure the proof of Theorem A.

2. Definitions and generic conditions

In this section, we present some definitions needed in later sections and generic conditions adopted as hypotheses in Theorem A.

2.1. **Definitions.**

Definition 2.1. Suppose that M is a 3-dimensional C^2 manifold. Let $\{l_\nu\}_{\nu\in J}$, $\{m_\nu\}_{\nu\in J}$ be C^2 families of regular curves in M, and let $\{S_\nu\}_{\nu\in J}$, $\{Y_\nu\}_{\nu\in J}$ be C^2 families of regular surfaces in M, where J is an open interval.

(1) Suppose that l_{ν_0} and m_{ν_0} intersect at a point s for some $\nu_0 \in J$ and some open neighborhood U of s in M has a C^2 change of coordinates with respect to which $m_\nu = \{(0,0,z) \in U\}$ for any $\nu \in J$ near ν_0. We say that s is a *quasi-transverse intersection* of l_{ν_0} and m_{ν_0} if

$$\dim(T_s(l_{\nu_0}) + T_s(m_{\nu_0})) = 2.$$

Moreover, s *unfolds generically* at $\nu = \nu_0$ with respect to $\{l_\nu\}_{\nu\in J}$, $\{m_\nu\}_{\nu\in J}$ if there exists a C^2 map $s : J \to M$ with $s(\nu) = s_\nu \in l_\nu$ for any $\nu \in J$ and $s(\nu_0) = s$ and a C^2 function $d : J \to \mathbb{R}^+$ with $d(\nu_0) \neq 0$ such that

$$T_sM = T_s(l_{\nu_0}) \oplus N \oplus T_s(m_{\nu_0}) \quad \text{and} \quad \mathrm{dist}(s_\nu, m_\nu) = |\nu - \nu_0| d(\nu) \tag{2.1}$$

for any ν near ν_0, where N is the one-dimensional space spanned by the non-zero tangent vector $(ds_\nu/d\nu)|_{\nu=\nu_0}$. This property corresponds to the conditions (GU1)–(GU3) in [9, §2.2.1].

(2) Suppose that l_{ν_0} and S_{ν_0} intersect at a point τ for some $\nu_0 \in J$. We say that τ is a *quadratic tangency* (or a *contact of order* 1) of l_{ν_0} and S_{ν_0} if there exists some C^2 change of coordinates on an open neighborhood $U(\tau)$ of τ such that $\tau = (0,0,0)$, $S_\nu = \{(x,y,z) \in U(\tau)\,;\, z = 0\}$ and l_ν has a regular parametrization $l(\nu,t) = (x(\nu,t), y(\nu,t), z(\nu,t))$ with $l(\nu_0, 0) = (0,0,0)$ and

$$\frac{\partial z}{\partial t}(\nu_0, 0) = 0 \quad \text{and} \quad \frac{\partial^2 z}{\partial t^2}(\nu_0, 0) \neq 0,$$

The tangency τ is said to *unfold generically* at $\nu = \nu_0$ with respect to $\{l_\nu\}_{\nu\in J}$ and $\{S_\nu\}_{\nu\in J}$ if

$$\frac{\partial z}{\partial \nu}(\nu_0, 0) \neq 0.$$

(3) Suppose that S_{ν_0} and Y_{ν_0} intersect at a point r for some $\nu_0 \in J$. We say that r is a *nondegenerate heterodimensional tangency* of S_{ν_0} and Y_{ν_0} if there exists a C^2 coordinate on an open set U in M containing r with $r = (u_0, v_0, 0)$ for some $u_0, v_0 \in \mathbb{R}$, $S_\nu = \{(x, y, z) \in U\,;\, z = 0\}$ and such that Y_ν has a parametrization $(x, y, f_\nu(x, y))$ the third entry $f_\nu(x, y) = f(\nu, x, y)$ of which is a C^2 function satisfying

$$f_{\nu_0}(u_0, v_0) = 0, \quad \frac{\partial f_{\nu_0}}{\partial x}(u_0, v_0) = \frac{\partial f_{\nu_0}}{\partial y}(u_0, v_0) = 0, \quad \det(Hf_{\nu_0})_{(u_0,v_0)} \neq 0, \tag{2.2}$$

where $(Hf_{\nu_0})_{(u_0,v_0)}$ is the Hessian matrix of f_{ν_0} at $(x, y) = (u_0, v_0)$.

The tangency r *unfolds generically* at $\nu = \nu_0$ if

$$\frac{\partial f}{\partial \nu}(\nu_0, u_0, v_0) \neq 0.$$

Remark 2.2. It is easy to see that the property (1) does not depend on the coordinates used to set l_ν in the z-axis. Similarly, the properties (2) and (3) do not depend on the coordinates used to set S_ν in the xy-plane.

When $\det(Hf_{\nu_0})_{(u_0,v_0)} > 0$ (resp. < 0) in Definition 2.1 (3), we say that the tangency $r = (u_0, v_0, 0)$ is of *elliptic* (resp. *hyperbolic*) type. The Taylor expansion of f_{ν_0} around (u_0, v_0) is

$$\begin{aligned} f_{\nu_0}(x, y) = {} & \frac{1}{2}\frac{\partial^2 f_{\nu_0}}{\partial x^2}(u_0, v_0)(x - u_0)^2 + \frac{\partial^2 f_{\nu_0}}{\partial x \partial y}(u_0, v_0)(x - u_0)(y - v_0) \\ & + \frac{1}{2}\frac{\partial^2 f_{\nu_0}}{\partial y^2}(u_0, v_0)(y - v_0)^2 + o\big((|x - u_0| + |y - v_0|)^2\big). \end{aligned} \tag{2.3}$$

From (2.3) together with the classification of quadratic surfaces in $\mathbb{R}^3$, we know that Y_{ν_0} has the form near $r = (u_0, v_0, 0)$ as illustrated in Fig. 2.1.



FIGURE 2.1. (1) r is of elliptic type and $\partial^2 f_{\nu_0}(u_0, v_0)/\partial x^2 < 0$. (2) r is of elliptic type and $\partial^2 f_{\nu_0}(u_0, v_0)/\partial x^2 > 0$. (3) r is of hyperbolic type.

2.2. **Generic conditions.** Throughout the remainder of this paper, we suppose that φ is a 3-dimensional C^2 diffeomorphism with saddle fixed points p of index$(p) = 1$ and q of index$(q) = 2$, and such that $W^s(p)$ and $W^u(q)$ have a nondegenerate heterodimensional tangency r, $W^u(p)$ and $W^s(q)$ have a quasi-transverse intersection s. The φ is *locally C^2 linearizable* in a neighborhood $U(q)$ of q if there exists a C^2 linearizing coordinate (x, y, z) on $U(q)$, that is,

$$q = (0, 0, 0), \quad \varphi(x, y, z) = (\alpha x, \beta y, \gamma z) \tag{2.4}$$

for any $(x, y, z) \in U(q)$ with $\varphi(x, y, z) \in U(q)$, where α, β and γ are eigenvalues of $(d\varphi)_q$.

One can take a local unstable manifold $W^u_{\mathrm{loc}}(q)$ so that it is an open disk in the plane $\{z = 0\}$ centered at $(x, y) = (0, 0)$. We may assume that the both points r, s are contained in $U(q)$ if necessary replacing r (resp. s) by $\varphi^{-n}(r)$ (resp. $\varphi^n(s)$) with sufficiently large $n \in \mathbb{N}$. We set

$$r = (u_0, v_0, 0)$$

with respect to the linearizing coordinate on $U(q)$.

We suppose moreover that $\{\varphi_{\mu,\nu}\}$ is a two-parameter family in $\mathrm{Diff}^2(M)$ with $\varphi_{0,0} = \varphi$ and satisfying the conditions of Theorem A. In particular, $\varphi_{\mu,\nu}$ is locally C^2 linearizable in a small neighborhood $U(q_{\mu,\nu})$ of $q_{\mu,\nu}$ in M and hence $\varphi_{\mu,\nu}$ has the form as (2.4) in $U(q_{\mu,\nu})$, where α, β, γ are C^2 functions on μ, ν, i.e., $\alpha = \alpha_{\mu,\nu}, \beta = \beta_{\mu,\nu}, \gamma = \gamma_{\mu,\nu}$.

We will put the following generic conditions (C1)-(C4) as the hypotheses in Theorem A.

(C1) (Generic condition for q) The φ is locally C^2 linearizable at q given as in (2.4). For simplicity, we suppose that every eigenvalues of $(d\varphi)_q$ is positive, that is,

$$0 < \gamma < 1 < \beta < \alpha.$$

(C2) (Generic unfolding property for r) The nondegenerate heterodimensional tangency r of $W^u(q)$ and $W^s(p)$ unfolds generically with respect to the μ-parameter families $\{W^u(q_{\mu,0})\}$ and $\{W^s(p_{\mu,0})\}$.

(C3) (Generic unfolding property for s) The quasi-transverse intersection s of $W^s(q)$ and $W^u(p)$ unfolds generically with respect to the ν-parameter families $\{W^s(q_{0,\nu})\}$ and $\{W^u(p_{0,\nu})\}$.

(C4) (Additional generic conditions) The tangency r is not on the x-axis $W^{uu}_{\mathrm{loc}}(q)$, that is,

$$v_0 \neq 0. \tag{2.5}$$

There exists a regular parametrization $l(t) = (x(t), y(t), z(t))$ $(t \in I)$ of a small curve in $W^u(p) \cap U(q)$ with respect to the linearizing coordinate (x, y, z) on $U(q)$ with $s = l(0)$ and

$$\frac{dx}{dt}(0) \neq 0, \tag{2.6}$$

where I is an open interval centered at 0.

There exists a C^2 function $f : O \to \mathbb{R}$ defined on an open disk O in the xy-plane centered at r such that $f(u_0, v_0) = 0$, $\{(x, y, f(x, y))\,;\, (x, y) \in O\} \subset W^s(p) \cap U(q)$ and

$$\frac{\partial^2 f}{\partial x^2}(u_0, v_0) \neq 0. \tag{2.7}$$

Note that the condition (2.7) is automatically satisfied when r is of elliptic type.

3. Some lemmas about parametrization and curvatures

The goal of this section is to prove three lemmas needed for the proof of Theorem A. These play important roles in Section 4.

- Lemma 3.1 presents a new parameter $(\hat{\mu}, \hat{\nu})$ such that, for any $\hat{\mu}$ near 0, there exists a quasi-transverse intersection $s_{\hat{\mu},0}$ of $W^s(q_{\hat{\mu},0})$ and $W^u(p_{\hat{\mu},0})$ which unfolds generically at $\hat{\nu} = 0$ with respect to the $\hat{\nu}$-parameter. After Lemma 3.1, we denote the new parameter $(\hat{\mu}, \hat{\nu})$ again by (μ, ν) for simplicity.
- In Lemma 3.2, we show that, for any μ_0 near 0, there exists a regular curve l_m in $W^u(p_{\mu_0,0})$ containing the quasi-transverse intersection $\varphi^m_{\mu_0,0}(s_{\mu_0,0})$ and arbitrarily C^2 close to $W^{uu}_{\mathrm{loc}}(q_{\mu_0,0})$. In particular, this implies that the curvature of l_m can be taken arbitrarily close to 0 with respect to the linearizing coordinate (2.4) on $U(q_{\mu_0,0})$.
- Lemma 3.3 gives a connection between the curvature and quadratic tangencies. In fact, we show that a tangency τ of a regular curve l and a regular surface S in $\mathbb{R}^3$ is quadratic if the curvature of l at τ is different from the normal curvature of S at τ along the direction tangent to l.

For any (μ, ν) near $(0, 0)$, we may assume that $U(q_{\mu,\nu})$ is equal to

$$D(\delta) := (-\delta, \delta)^3$$

with respect to the linearizing coordinate given in Subsection 2.2 for some constant $\delta > 0$. Since s is a quasi-transverse intersection which unfolds generically with respect to the ν-parameter families $\{W^s(q_{0,\nu})\}$ and $\{W^u(p_{0,\nu})\}$ by the condition (C3), there exists a C^2 continuation $\hat{s}_\nu \in W^u(p_{0,\nu}) \cap D(\delta)$ with $\hat{s}_0 = s$ and such that $\hat{s}_\nu$ satisfies the conditions same as those for s_ν in Definition 2.1 (1). By (2.6), for any ν near 0, the component l_ν of $W^u(p_{0,\nu}) \cap D(\delta)$ containing $\hat{s}_\nu$ meets transversely the yz-plane at a point s_ν which defines a C^2 continuations $\{s_\nu\}$ with $s_0 = s$, see Fig. 3.1. Note that $d\hat{s}_\nu/d\nu(0) = ds_\nu/d\nu(0) + \boldsymbol{w}$ for some $\boldsymbol{w} \in T_s(l_0) = T_s(W^u(p))$, where



Figure 3.1.

$d\hat{s}_\nu/d\nu(0)$ denotes $d\hat{s}_\nu/d\nu|_{\nu=0}$. Let $y(\nu)$ be the y-coordinate of s_ν. If $dy/d\nu(0) = 0$, then $ds_\nu/d\nu(0)$ would be tangent to the z-axis $W^s_{\mathrm{loc}}(q)$ at s and hence $d\hat{s}_\nu/d\nu(0) \in T_s(W^s(q)) \oplus T_s(W^u(p))$. This contradicts (2.1). Thus, we have

$$\frac{dy}{d\nu}(0) \neq 0. \tag{3.1}$$

For any (μ, ν) near $(0,0)$, let $s_{\mu,\nu}$ be a transverse intersection point of $W^u(p_{\nu,\nu}) \cap D(\delta)$ with the yz-plane such that $\{s_{\mu,\nu}\}$ is a C^2 continuation with $s_{0,\nu} = s_\nu$.

Lemma 3.1. *There exists a constant $\rho > 0$ and a C^2 function $\tilde{\nu} : (-\rho, \rho) \to \mathbb{R}$ such that, for any $\mu \in (-\rho, \rho)$, $s_{\mu,\tilde{\nu}(\mu)}$ is a quasi-transverse intersection of $W^s(q_{\mu,\tilde{\nu}(\mu)})$ and $W^u(p_{\mu,\tilde{\nu}(\mu)})$ which unfolds generically with respect to the ν-parameter families $\{W^s(q_{\mu(\mathrm{fixed}),\nu})\}$ and $\{W^u(p_{\mu(\mathrm{fixed}),\nu})\}$.*

Proof. Let $y(\mu, \nu)$ be the y-coordinate of $s_{\mu,\nu}$. By (3.1), $\partial y/\partial \nu(0,0) \neq 0$. Hence, by the Implicit Function Theorem, there exists a C^2 function $\tilde{\nu} : (-\rho, \rho) \to \mathbb{R}$ for some $\rho > 0$ such that

$$\tilde{\nu}(0) = 0, \ y(\mu, \tilde{\nu}(\mu)) = 0, \ \frac{\partial y}{\partial \nu}(\mu, \tilde{\nu}(\mu)) \neq 0$$

for any $\mu \in (-\rho, \rho)$. This implies that $s_{\mu,\tilde{\nu}(\mu)}$ is a quasi-transverse intersection unfolding generically at $\nu = \tilde{\nu}(\mu)$ with respect to the ν-parameter families $\{W^s(q_{\mu,\nu})\}$ and $\{W^u(p_{\mu,\nu})\}$. □

A new parametrization. Consider the coordinate $(\hat{\mu}, \hat{\nu})$ on the parameter space defined by $\hat{\mu} = \mu, \hat{\nu} = \nu - \tilde{\nu}(\mu)$. For simplicity, we denote the new coordinate again by (μ, ν). Thus, there exists a continuation $\{s_{\mu,0}\}_{\mu \in (-\rho,\rho)}$ of quasi-transverse intersections of $W^s(q_{\mu,0})$ and $W^u(p_{\mu,0})$ such that each $s_{\mu,0}$ unfolds generically at $\nu = 0$ with respect to the ν-parameter families $\{W^s(q_{\mu,\nu})\}$ and $\{W^u(p_{\mu,\nu})\}$.

Fix μ_0 with sufficiently small $|\mu_0|$ arbitrarily. By the properties (2.4) and (2.6), there exists $m_0 \in \mathbb{N}$ such that, for any $m \geq m_0$, one can parameterize the component l_m of $W^u(p_{\mu_0,0}) \cap D(\delta)$ containing $\varphi^m_{\mu_0,0}(s_{\mu_0,0})$ so that $l_m(0) = \varphi^m_{\mu_0,0}(s_{\mu_0,0})$ and

$$l_m(t) = (t, y_m(t), z_m(t)) \quad (t \in (-\delta, \delta)).$$

Lemma 3.2. *The sequence $\{l_m\}$ C^2 converges uniformly to $W^{uu}_{\mathrm{loc}}(q_{\mu_0,0})$ as $m \to \infty$. In particular, for any $\varepsilon > 0$, there exists $\hat{m}_0 \geq m_0$ such that the curvature at any point of l_m is less than ε with respect to the standard Euclidean metric on $U(q_{\mu_0,0}) = D(\delta)$ if $m \geq \hat{m}_0$.*

Proof. By (2.4), for any $m \geq m_0$,

$$l_m(t) = (t, \beta^n y_{m_0}(\alpha^{-n} t), \gamma^n z_{m_0}(\alpha^{-n} t)),$$

where $n = m - m_0$, $\alpha = \alpha_{\mu_0,0}$, $\beta = \beta_{\mu_0,0}$, $\gamma = \gamma_{\mu_0,0}$. Thus we have

$$\begin{aligned} \frac{dl_m}{dt}(t) &= \left(1, \ \frac{\beta^n}{\alpha^n}\frac{dy_{m_0}}{dt}(\alpha^{-n}t), \ \frac{\gamma^n}{\alpha^n}\frac{dz_{m_0}}{dt}(\alpha^{-n}t)\right) \xrightarrow{\text{uniformly}} (1,0,0), \\ \frac{d^2 l_m}{dt^2}(t) &= \left(0, \ \frac{\beta^n}{\alpha^{2n}}\frac{d^2 y_{m_0}}{dt^2}(\alpha^{-n}t), \ \frac{\gamma^n}{\alpha^{2n}}\frac{d^2 z_{m_0}}{dt^2}(\alpha^{-n}t)\right) \xrightarrow{\text{uniformly}} (0,0,0) \end{aligned} \tag{3.2}$$

as $m \to \infty$. Since $\{l_m(0)\}_{m=m_0}^{\infty}$ converges to $q_{\mu_0,0} = (0,0,0)$, it follows from (3.2) that $\{l_m\}$ C^2 converges uniformly to the x-axis in $D(\delta)$. □

Lemma 3.3. *Let S be a regular surface in the Euclidean 3-space $\mathbb{R}^3$ and l a regular curve tangent to S at τ. Suppose that the curvature $\kappa_l(\tau)$ of l at τ is less than the absolute value of the normal curvature $\kappa_S(\tau, \boldsymbol{w})$ of S at τ along a non-zero vector $\boldsymbol{w}$ tangent to l. Then tangency of S and l at τ is quadratic.*

Proof. By changing the coordinate (x, y, z) on $\mathbb{R}^3$ by an isometry, we may assume that $\tau = (0, 0, 0)$, the tangent space of S at τ is the xy-plane and $\boldsymbol{w}/\|\boldsymbol{w}\| = (1, 0, 0)$. Then one can suppose that S (resp. l) is parameterized as $(x, y, \psi(x, y))$ (resp. $(x, f_1(x), f_2(x))$ in a small neighborhood of $(0, 0, 0)$ in $\mathbb{R}^3$. Since the graph of $z = \psi(x, 0)$ is the cross section of S along the xz-plane,

$$|\kappa_S(\tau, \boldsymbol{w})| = \frac{|g''(0)|}{(g'(0)^2 + 1)^{3/2}} = |g''(0)|,$$

where $g(x) = \psi(x, 0)$. Since the graph of $z = f_2(x)$ coincides with the orthogonal projection $\bar{l}$ of l into the xz-plane,

$$\kappa_l(\tau) \geq \kappa_{\bar{l}}(\tau) = \frac{|f_2''(0)|}{(f_2'(0)^2 + 1)^{3/2}} = |f_2''(0)|.$$

It follows from our assumption $|\kappa_S(\tau, \boldsymbol{w})| > \kappa_l(\tau)$ that $|g''(0)| > |f_2''(0)|$. This shows that the tangency at τ is quadratic. $\square$

4. Proof of Theorem A

In this section, we give the proof of Theorem A.

- In Subsection 4.1, we show that, for any μ_0 in either $(-\varepsilon, 0)$ or $(0, \varepsilon)$ and any sufficiently large $m \in \mathbb{N}$, there exists ν_m with $\lim_{m\to\infty} \nu_m = 0$ such that $W^u(p_{\mu_0,\nu_m})$ and $W^s(p_{\mu_0,\nu_m})$ have a quadratic tangency τ_m (Assertion 4.2 and Assertion 4.3). Here the sign of μ_0 is chosen so that $\mu_0 \cdot b_{\mu_0,0} < 0$ (resp. $\mu_0 \cdot b_{\mu_0,0} > 0$) if the tangency r is of elliptic (resp. hyperbolic) type, where $b_{\mu_0,0}$ is the coefficient of $(x - \nu_{\mu_0,0})^2$-term of the Taylor expansion (4.1). See Fig. 4.1 and Fig. 4.4. As is suggested in Fig. 4.2, the existence of the homoclinic tangency τ_m is proved by using the Intermediate Value Theorem. By Lemma 3.2, the curvature of $W^u(p_{\mu_0,\nu_m})$ at τ_m converges to zero as $m \to \infty$. On the other hand, we will show that the normal curvature of $W^s(p_{\mu_0,\nu_m})$ at τ_m along the direction tangent to l_m is bounded away from zero. Hence, by Lemma 3.3, the tangency τ_m is quadratic.
- In Subsection 4.2, we show that the quadratic homoclinic tangency τ_m unfolds generically at $\nu = \nu_m$ with respect to the ν-parameter families $\{W^s(p_{\mu_0,\nu})\}$ and $\{W^u(p_{\mu_0,\nu})\}$ by representing a neighborhood of τ_m in $W^s(p_{\mu_0,\nu})$ as the graph of a function of (x, z).

4.1. **Existence of quadratic homoclinic tangencies.** Let $\{\varphi_{\mu,\nu}\}$ be the family given in Subsection 2.2. In particular, $r = (u_0, v_0, 0)$ is a nondegenerate heterodimensional tangency of $W^u(q)$ and $W^s(p)$ which unfolds generically with respect to the μ-parameter families $\{W^u(q_{\mu,0})\}$ and $\{W^s(p_{\mu,0})\}$. By our settings in Sections 2 and 3, there exist C^2 functions $f_{\mu,\nu} : O \subset \mathbb{R}^2 \to \mathbb{R}$ C^2 depending on (μ, ν) with $f_{0,0} = f$ and

$$\Sigma(\mu, \nu) := \{(x, y, f_{\mu,\nu}(x, y))\,;\, (x, y) \in O\} \subset W^s(p_{\mu,\nu}) \cap D(\delta)$$

for any (μ, ν) near $(0, 0)$. Since $\det(Hf)_{(u_0,v_0)} \neq 0$, there exists a uniquely determined C^2 continuation $(u_{\mu,\nu}, v_{\mu,\nu})$ with $(u_{0,0}, v_{0,0}) = (u_0, v_0)$ and

$$\frac{\partial f_{\mu,\nu}}{\partial x}(u_{\mu,\nu}, v_{\mu,\nu}) = \frac{\partial f_{\mu,\nu}}{\partial y}(u_{\mu,\nu}, v_{\mu,\nu}) = 0.$$

Proposition 4.1. *For a sufficiently small $\varepsilon > 0$ and any μ in either $(0,\varepsilon)$ or $(-\varepsilon,0)$, there exists ν arbitrarily close to 0 such that $\varphi_{\mu,\nu}$ has a quadratic homoclinic tangency associated to $p_{\mu,\nu}$.*

By the condition (2.5), r is not in the x-axis. One can take the linearizing coordinate on $D(\delta)$ so that s (resp. r) is in the upper half space $\{z > 0\}$ (resp. $\{x > 0\}$). The Taylor expansion of $f_{\mu,\nu}$ around $(u_{\mu,\nu}, v_{\mu,\nu})$ has the form:

$$\begin{aligned} f_{\mu,\nu}(x,y) = a_{\mu,\nu} &+ \frac{1}{2}b_{\mu,\nu}(x-u_{\mu,\nu})^2 + c_{\mu,\nu}(x-u_{\mu,\nu})(y-v_{\mu,\nu}) \\ &+ \frac{1}{2}d_{\mu,\nu}(y-v_{\mu,\nu})^2 + o\big((|x-u_{\mu,\nu}|+|y-v_{\mu,\nu}|)^2\big), \end{aligned} \tag{4.1}$$

where $a_{0,0} = 0$ and

$$b_{\mu,\nu} = \frac{\partial^2 f_{\mu,\nu}}{\partial x^2}(u_{\mu,\nu}, v_{\mu,\nu}),\ c_{\mu,\nu} = \frac{\partial^2 f_{\mu,\nu}}{\partial x \partial y}(u_{\mu,\nu}, v_{\mu,\nu}),\ d_{\mu,\nu} = \frac{\partial^2 f_{\mu,\nu}}{\partial y^2}(u_{\mu,\nu}, v_{\mu,\nu}).$$

Since the tangency r unfolds generically with respect to $\varphi = \varphi_{0,0}$ by (C1),

$$\eta_0 = \frac{\partial a_{\mu,\nu}}{\partial \mu}\Big|_{(\mu,\nu)=(0,0)} \neq 0. \tag{4.2}$$

If necessary replacing μ by $-\mu$, we may assume that $\eta_0 > 0$. By the condition (2.7), $b_{0,0} \neq 0$ and hence $b_{\mu,\nu} \neq 0$ for any (μ,ν) near $(0,0)$.

Proposition 4.1 is divided to the following two assertions.

Assertion 4.2 (Elliptic case)**.** *If r is of elliptic type, then Proposition 4.1 holds.*

Proof. First we consider the case of $b_{\mu,\nu} < 0$ for any (μ,ν) near $(0,0)$. By (4.2), for any sufficiently small $\mu_0 > 0$, the intersection $C_{\mu_0} = \Sigma(\mu_0, 0) \cap \{z = 0\}$ is a circle disjoint from the x-axis. For a sufficiently small $h_0 > 0$, $A = \Sigma(\mu_0,0) \cap \{0 \le z \le h_0\}$ is an annulus in $D(\delta)$, see Fig. 4.1 (1). Replacing m_0 by an integer greater than m_0 if necessary, we may assume that $z_m(0) < h_0/2$ for any $m \geq m_0$. By Lemma 3.2, the curve $l_m \subset W^u(p_{\mu_0,0}) \cap D(\delta)$ given in Section 3 is sufficiently C^2 close to the x-axis. Thus one can suppose that $\pi_y(l_m) \cap \pi_y(A) = \emptyset$, where $\pi_y : D(\delta) \longrightarrow \mathbb{R}$ is



Figure 4.1. (1) The case of $\mu_0 > 0, \nu = 0, b_{\mu_0,0} < 0$. (2) The case of $\mu_0 < 0, \nu = 0, b_{\mu_0,0} > 0$. Each shaded region represents A.

the orthogonal projection defined by $\pi_y(x,y,z)=y$. For any sufficiently small ν, let $l_{m,\nu}$ be the component of $W^u(p_{\mu_0,\nu})\cap D(\delta)$ such that $\{l_{m,\nu}\}$ is an ν-continuation with $l_{m,0}=l_m$, and set $A_\nu=\Sigma(\mu_0,\nu)\cap\{0\le z\le h_0\}$. Moreover, one can suppose that $l_{m,\nu}$ is parameterized as

$$l_{m,\nu}(t)=(t,y_m(\nu,t),z_m(\nu,t))\quad (t\in(-\delta,\delta)).$$

By the condition (C3), one can take $\bar\nu\neq 0$ with arbitrarily small $|\bar\nu|$ such that

$$0<\pi_y(l_{m_0,\bar\nu}(0))\le \sup\{\pi_y(l_{m_0,\bar\nu})\}<\min\{\pi_y(A_{\bar\nu})\}.$$

We may assume that $\bar\nu>0$ if necessary replacing ν by $-\nu$. For any integer m sufficiently greater than m_0, there exists $0<\bar\nu_m<\bar\nu$ such the continuation $\{l_{m,\nu}\}_{0\le\nu\le\bar\nu_m}$ is well defined and

$$\max\{\pi_y(A_{\bar\nu_m})\}<\inf\{\pi_y(l_{m,\bar\nu_m})\}$$

holds, see Fig. 4.2 (1). By the Intermediate Value Theorem, there exists $0<\nu_m<\bar\nu_m$ such that l_{m,ν_m} and A_{ν_m} have a tangency τ_m, see Fig. 4.2 (2). Since $l_{m,\nu_m}\subset W^u(p_{\mu_0,\nu_m})$ and $A_{\nu_m}\subset W^s(p_{\mu_0,\nu_m})$, τ_m is a homoclinic tangency associated to p_{μ_0,ν_m}.



(1) (2)

Figure 4.2. The cross sections.

When $b_{\mu,\nu}>0$ for any (μ,ν) near $(0,0)$, one can prove the existence of a homoclinic tangency τ_m near r associated to p_{μ_0,ν_m} by arguments quite similar to those as above for any μ_0 with $\mu_0<0$.

It remains to show that the tangency τ_m is quadratic. Since $\Sigma(\mu_0,\nu_m)$ is of elliptic type and $\lim_{m\to\infty}\nu_m=0$, any normal curvature of $\Sigma(\mu_0,\nu_m)$ at τ_m is greater than some positive constant κ_0 independent of m. On the other hand, by an argument quite similar to that in Lemma 3.2, for any m sufficiently greater than m_0, the curvature of l_{m,ν_m} at τ_m is less than κ_0. Thus, by Lemma 3.3, τ_m is a quadratic tangency. □

Assertion 4.3 (Hyperbolic case)**.** *When r is a tangency of hyperbolic type, Proposition 4.1 holds.*

Proof. Since r is of hyperbolic type, $\Sigma(0,0)\cap\{z=0\}$ consists of two almost straight curves α_1,α_2 meeting transversely at r, see Fig. 4.3 (1). If necessary by reducing the domain O of $f_{\mu,\nu}$ containing $(u_{\mu,\nu},v_{\mu,\nu})$, we may assume that $\Sigma(\mu,\nu)\cap\{z=0\}$



FIGURE 4.3. (1) The situation when $(\mu,\nu)=(0,0)$. (2) The situation when $(\mu,\nu)=(\mu_0,0)$.

is disjoint from the x-axis for any (μ,ν) near $(0,0)$. If $\boldsymbol{w}_i=(\xi_i,\eta_i,0)$ $(i=1,2)$ is a unit vector tangent to α_i at r, then $b_{0,0}\xi_i^2+2c_{0,0}\xi_i\eta_i+d_{0,0}\eta_i^2=0$. This implies that the normal curvature $\kappa_{\Sigma(0,0)}(r,\boldsymbol{w}_i)$ of $\Sigma(0,0)$ at r along $\boldsymbol{w}_i$ is zero. Since $b_{0,0}\neq 0$ by (2.7), both $\boldsymbol{w}_1$, $\boldsymbol{w}_2$ are not parallel to the unit tangent vector $\boldsymbol{v}_0=(1,0,0)$. Thus we have $\kappa_{\Sigma(0,0)}(r,\boldsymbol{v}_0)\neq 0$. When $b_{0,0}<0$ (resp. $b_{0,0}>0$), for any sufficiently small μ_0 with $\mu_0<0$ (resp. $\mu_0>0$), $\Sigma(\mu_0,0)\cap\{z=0\}$ consists of two C^2 curves β_1,β_2 separated by a line in the xy-plane parallel to x-axis, see Fig. 4.3 (2), and

$$\kappa_0:=|\kappa_{\Sigma(\mu_0,0)}(\tau,\boldsymbol{v}_0)|>0, \tag{4.3}$$

where τ is a point of $\beta_1\cup\beta_2$ the tangent line at which is parallel to the x-axis. One can take $\bar\nu>0$ and $h_0>0$ so that $A_\nu=\Sigma(\mu_0,\nu)\cap\{0\leq z\leq h_0\}$ is a disjoint union of two curvilinear rectangles for any ν with $0\leq\nu\leq\bar\nu$, see Fig. 4.4. Moreover,



FIGURE 4.4. (1) The case of $\mu_0<0,\nu=0,b_{\mu_0,0}<0$. (2) The case of $\mu_0>0,\nu=0,b_{\mu_0,0}>0$. Each pair of the shaded regions represents A.

by (4.3), the $\bar\nu>0$ can be chosen so that $|\kappa_{\Sigma(\mu_0,\nu)}(\tilde\tau,\boldsymbol{w})|>\kappa_0/2$ for any point $\tilde\tau\in A_{\mu_0,\nu}$ sufficiently near τ and any unit vector $\boldsymbol{w}\in T_{\tilde\tau}(A_{\mu_0,\nu})$ sufficiently near $\boldsymbol{v}_0$. As in the proof of Assertion 4.2, for any integer m sufficiently greater than

m_0, there exists ν_m with $0 < \nu_m < \bar{\nu}$ and such that $l_{m,\nu_m} \subset W^s(p_{\mu_0,\nu_m})$ and $A_{\nu_m} \subset W^u(p_{\mu_0,\nu_m})$ have a quadratic tangency τ_m, see Fig. 4.5. $\square$



(1) (2)

Figure 4.5.

4.2. **Generic unfolding of the tangency.** For short, set $p_{\mu_0,\nu} = p_\nu$, $f_{\mu_0,\nu}(x,y) = f_\nu(x,y)$ and $(u_{\mu_0,\nu}, v_{\mu_0,\nu}) = (u_\nu, v_\nu)$.

Let $\tau_m = (\hat{x}_m, \hat{y}_m, f_{\nu_m}(\hat{x}_m, \hat{y}_m))$ be the homoclinic tangency of $W^u(p_{\nu_m})$ and $W^s(p_{\nu_m})$ given in Proposition 4.1. From (4.1), we have

$$\frac{\partial f_{\nu_m}}{\partial y}(x,y) = c_m(x - u_{\nu_m}) + d_m(y - v_{\nu_m}) + o_1,$$
$$\frac{\partial f_{\nu_m}}{\partial x}(x,y) = b_m(x - u_{\nu_m}) + c_m(y - v_{\nu_m}) + o_1,$$

where $b_m = b_{\mu_0,\nu_m}$, $c_m = c_{\mu_0,\nu_m}$, $d_m = d_{\mu_0,\nu_m}$ and $o_1 = o(|x - u_{\nu_m}| + |y - v_{\nu_m}|)$. Thus $b_m \partial f_{\nu_m}(x,y)/\partial y - c_m \partial f_{\nu_m}(x,y)/\partial x = (b_m d_m - c_m^2)(y - v_{\nu_m}) + o_1$. On the other hand, since there exists a unit vector tangent to $\Sigma(\mu_0, \nu_m)$ at τ_m converges to $(1,0,0)$ as $m \to \infty$, $\lim_{m\to\infty} \partial f_{\nu_m}(\hat{x}_m, \hat{y}_m)/\partial x = 0$. Since $\lim_{m\to\infty} b_m = b_{\mu_0,0} \neq 0$ and $\lim_{m\to\infty} b_m d_m - c_m^2 = \det(Hf_{\mu_0,0})(u_{\mu_0,0}, v_{\mu_0,0}) \neq 0$,

$$\frac{\partial f_{\nu_m}}{\partial y}(\hat{x}_m, \hat{y}_m) = \frac{c_m}{b_m}\frac{\partial f_{\nu_m}}{\partial x}(\hat{x}_m, \hat{y}_m) + \frac{b_m d_m - c_m^2}{b_m}(\hat{y}_m - v_{\nu_m}) + o_1 \neq 0$$

for all sufficiently large m. By the Implicit Function Theorem, there exists a C^2 function $y = g_\nu(x,z) = g(\nu,x,z)$ defined in a small neighborhood of $(\nu_m, \hat{x}_m, f_{\nu_m}(\hat{x}_m, \hat{y}_m))$ in the (ν,x,z)-space with

$$(x, y, f_\nu(x,y)) = (x, g_\nu(x,z), z).$$

Proposition 4.4. *For all sufficiently large m, the quadratic homoclinic tangency τ_m of $W^s(p_{\nu_m})$ and $W^u(p_{\nu_m})$ unfolds generically at $\nu = \nu_m$ with respect to the ν-parameter families $\{W^s(p_\nu)\}$ and $\{W^u(p_\nu)\}$.*

Proof. Recall that $l_{m,\nu}$ has the parametrization $l_{m,\nu}(t) = (t, y_m(\nu,t), z_m(\nu,t))$ with $l_{m,\nu_m}(\hat{x}_m) = \tau_m$. By Definition 2.1 (2), it suffices to show that

$$\frac{\partial y_m}{\partial \nu}(\nu_m, \hat{x}_m) \neq \frac{\partial g}{\partial \nu}(\nu_m, \hat{x}_m, z_m(\nu_m, \hat{x}_m)) \tag{4.4}$$

for all sufficiently large m. Note that

$$\lim_{m\to\infty}\frac{\partial g}{\partial\nu}(\nu_m,\hat{x}_m,z_m(\nu_m,\hat{x}_m))=\frac{\partial g}{\partial\nu}(0,\hat{x}_\infty,0),$$

where $\hat{x}_\infty$ is the x-coordinate of a point τ in $\Sigma(\mu_0,0)\cap\{z=0\}$ the tangent line in xy-plane at which is parallel to $(1,0,0)$, see Fig. 4.3 (2) in the case that r is of hyperbolic type. If we set $\tilde{x}_{m,\nu}=\alpha_\nu^{-n}\hat{x}_m$, then $\varphi_\nu^n(l_{m_0,\nu}(\tilde{x}_{m,\nu}))=l_{m,\nu}(\hat{x}_m)$, where $n=m-m_0$ and $\alpha_\nu=\alpha_{\mu_0,\nu}$. As was seen in the proof of Lemma 3.1,

$$\lim_{m\to\infty}\frac{\partial y_{m_0}}{\partial\nu}(\nu_m,\tilde{x}_{m,\nu_m})=\frac{\partial y_{m_0}}{\partial\nu}(0,0)\neq 0. \tag{4.5}$$

We denote the ν-function $y_{m_0}(\nu,\tilde{x}_{m,\nu})$ by $h_m(\nu)$. Since $\lim_{m\to\infty}d\tilde{x}_{m,\nu}/d\nu=0$, it follows from (4.5) that

$$\begin{aligned}\left|\frac{dh_m}{d\nu}(\nu_m)\right|&=\left|\frac{\partial y_{m_0}}{\partial\nu}(\nu_m,\tilde{x}_{m,\nu_m})+\frac{\partial y_{m_0}}{\partial x}(\nu_m,\tilde{x}_{m,\nu_m})\frac{d\tilde{x}_{m,\nu}}{d\nu}(\nu_m)\right|\\&\geq\left|\frac{\partial y_{m_0}}{\partial\nu}(\nu_m,\tilde{x}_{m,\nu_m})\right|-\left|\frac{\partial y_{m_0}}{\partial x}(\nu_m,\tilde{x}_{m,\nu_m})\frac{d\tilde{x}_{m,\nu}}{d\nu}(\nu_m)\right|>C_0\end{aligned}\tag{4.6}$$

for some positive constant C_0 and all m sufficiently greater than m_0. Since $y_m(\nu,\hat{x}_m)=\beta_\nu^n h_m(\nu)$ for $\beta_\nu:=\beta_{\mu_0,\nu}$,

$$\begin{aligned}\frac{\partial y_m}{\partial\nu}(\nu_m,\hat{x}_m)&=\beta_{\nu_m}^n\frac{dh_m}{d\nu}(\nu_m)+n\beta_{\nu_m}^{n-1}\frac{d\beta_\nu}{d\nu}(\nu_m)h_m(\nu_m)\\&=\beta_{\nu_m}^n\frac{dh_m}{d\nu}(\nu_m)+\frac{n}{\beta_{\nu_m}}\frac{d\beta_\nu}{d\nu}(\nu_m)y_m(\nu_m,\hat{x}_m).\end{aligned}$$

Since $\lim_{m\to\infty}\beta_{\nu_m}=\beta_0>1$ and $|y_m(\nu_m,\hat{x}_m)|\leq\delta$, the inequality (4.6) implies $\lim_{m\to\infty}|\partial y_m(\nu_m,\hat{x}_m)/\partial\nu|=\infty$. This shows (4.4). □

Proof of Theorem A. Propositions 4.1 and 4.4 imply Theorem A. □

Acknowledgments

We would like to thank Bau-Sen Du and Yi-Chiuan Chen for their support and hospitality, and Ming-Chia Li and Mikhail Malkin for their discussions during the first draft of this paper was written in Academia Sinica of Taiwan. We also would like to thank the referees for their valuable comments.

References

[1] M. Asaoka, A simple construction of C^1-Newhouse domains for higher dimensions, *Proc. Amer. Math. Soc.*, **136** (2008), no. 2, 677–686.

[2] Ch. Bonatti and L. J. Díaz, Connexions hétéroclines et généricité d'une infinité de puits et de sources, *Ann. Sci. École Norm. Sup.*, **(4) 32** (1999), no. 1, 135–150.

[3] Ch. Bonatti, L. J. Díaz, and E. R. Pujals, A C^1-generic dichotomy for diffeomorphisms: Weak forms of hyperbolicity or infinitely many sinks or sources, *Ann. of Math.*, **158** (2003), 355–418.

[4] Ch. Bonatti and L. J. Díaz, Robust heterodimensional cycles and C^1-generic dynamics, *Journal of the Inst. of Math. Jussieu*, **7** 3 (2008), 469–525.

[5] L. J. Díaz, Robust nonhyperbolic dynamics and heterodimensional cycles, *Ergod. Th. Dynam. Sys.*, **15**(1995), 291–315.

[6] L. J. Diaz, A. Nogueira and E. R. Pujals, Heterodimensional tangencies, *Nonlinearity*, **19** (2006), 2543–2566.

[7] L. J. Díaz and J. Rocha, Non-connected heterodimensional cycles: bifurcation and stability, *Nonlinearity*, **5** (1992), 1315–1341.

[8] L. J. Díaz and J. Rocha, Large measure of hyperbolic dynamics when unfolding heteroclinic cycles, *Nonlinearity*, **10** (1997), 857–884.
[9] L. J. Díaz and J. Rocha, Partially hyperbolic and transitive dynamics generated by heteroclinic cycles, *Ergod. Th. Dynam. Sys.*, **21** (2001), 25–76.
[10] S. V. Gonchenko, D. V. Turaev, and L. P. Shilnikov, On the existence of Newhouse regions near systems with non-rough Poincaré homoclinic curve (multidimensional case) *Russ. Acad. Sci. Dokl. Math.*, **47** (1993), 268–273.
[11] S. V. Gonchenko, L. P. Shilnikov and D. V. Turaev, Dynamical phenomena in systems with structurally unstable Poincaré homoclinic orbits, *Chaos*, **6** (1996), 15–31.
[12] B. Leal, High dimension diffeomorphisms exhibiting infinitely many strange attractors, *Ann. I. H. Poincaré*, **25** (2008), 587–607.
[13] S. E. Newhouse, The abundance of wild hyperbolic sets and non-smooth stable sets for diffeomorphisms, *Publ. Math. I.H.É.S.*, **50** (1979), 101–151.
[14] J. Palis, A global perspective for non-conservative dynamics, *Ann. I. H. Poincaré*, **22** (2005), 485–507.
[15] J. Palis and M. Viana, High dimension diffeomorphisms displaying infinitely many periodic attractors, *Ann. of Math. (2)*, **140** (1994), 207–250.
[16] C. Robinson, Bifurcation to infinitely many sinks, *Commun. Math. Phys.*, **90** (1983), 433–459.
[17] N. Romero, Persistence of homoclinic tangencies in higher dimensions, *Ergod. Th. Dynam. Sys.*, **15** (1995), no. 4, 735–757.
[18] M. Viana, Strange attractors in higher dimensions, *Bol. Soc. Brasil. Mat. (N.S.)* **24** (1993), 13–62.

Department of Mathematics, Kyoto University of Education, 1 Fukakusa-Fujinomori, Fushimi-ku, Kyoto, 612-8522, Japan
E-mail address: `skiriki@kyokyo-u.ac.jp`

Department of Mathematics and Information Sciences, Tokyo Metropolitan University, Minami-Ohsawa 1-1, Hachioji, Tokyo 192-0397, Japan
E-mail address: `nishizawa-yusuke@ed.tmu.ac.jp`

Department of Mathematics and Information Sciences, Tokyo Metropolitan University, Minami-Ohsawa 1-1, Hachioji, Tokyo 192-0397, Japan
E-mail address: `tsoma@tmu.ac.jp`